

$B(0, N)$ -graded Lie superalgebras coordinatized by quantum tori

Hongjia Chen, Yun Gao*, Shikui Shang

Abstract

We use a fermionic extension of the bosonic module to obtain a class of $B(0, N)$ -graded Lie superalgebras with nontrivial central extensions.

0 Introduction

$B(M - 1, N)$ -graded Lie superalgebras were first investigated and classified up to central extension by Benkart-Elduque (see also Garcia-Neher's work in [GN]). Those root graded Lie superalgebras are a super-analog of root graded Lie algebras. Fermionic and bosonic representations for the affine Kac-Moody Lie algebras were studied by Frenkel [F1,2] and Kac-Peterson [KP]. Feingold-Frenkel [FF] constructed representations for all classical affine Lie algebras by using Clifford or Weyl algebras with infinitely many generators. They also obtained realizations for certain affine Lie superalgebras including the affine $B(0, N)$. [G] gave bosonic and fermionic representations for the extended affine Lie algebra $\widetilde{gl_N(\mathbb{C}_q)}$, where \mathbb{C}_q is the quantum torus in two variables. [CG] constructed modules for some BC_N -graded Lie algebras by considering a fermionic extension of the fermionic module.

In this paper, we will consider a fermionic extension of the bosonic module to obtain a class of $B(0, N)$ -graded Lie superalgebras with nontrivial central extensions.

*Research was partially supported by NSERC of Canada and Chinese Academy of Science. This paper is dedicated to Professor Sheng Gong on the occasion of his 75th birthday.

The organization of the paper is as follows. In Section 1, we review some basics on the quantum torus and present examples of $B(0, N)$ -graded Lie superalgebras coordinatized by quantum tori which are subalgebras of $\widehat{gl(1, 2N)}(\mathbb{C}_q)$. In Section 2, we use bosons and fermions to construct representations for those examples of $B(0, N)$ -graded Lie superalgebras.

Throughout this paper, we denote the field of complex numbers and the ring of integers by \mathbb{C} and \mathbb{Z} respectively.

1 $B(0, N)$ -graded Lie superalgebras

We first recall some basics on quantum tori and then go on to present examples of $B(0, N)$ -graded Lie superalgebras coordinatized by quantum tori. For more information on Lie superalgebras graded by root systems, see [BE1]-[BE2] and [GN].

Let q be a non-zero complex number. A quantum torus associated to q (see [M]) is the unital associative \mathbb{C} -algebra $\mathbb{C}_q[x^\pm, y^\pm]$ (or, simply \mathbb{C}_q) with generators x^\pm, y^\pm and relations

$$(1.1) \quad xx^{-1} = x^{-1}x = yy^{-1} = y^{-1}y = 1 \quad \text{and} \quad yx = qxy.$$

Set $\Lambda(q) = \{n \in \mathbb{Z} | q^n = 1\}$. From [BGK] we see that $[\mathbb{C}_q, \mathbb{C}_q]$ has a basis consisting of monomials $x^m y^n$ for $m \notin \Lambda(q)$ or $n \notin \Lambda(q)$.

Let $\bar{}$ be the anti-involution on \mathbb{C}_q given by

$$(1.2) \quad \bar{x} = x, \quad \bar{y} = y^{-1}.$$

We have $\mathbb{C}_q = \mathbb{C}_q^+ \oplus \mathbb{C}_q^-$, where $\mathbb{C}_q^\pm = \{s \in \mathbb{C}_q | \bar{s} = \pm s\}$, then

$$(1.3) \quad \begin{aligned} \mathbb{C}_q^+ &= \text{span}\{x^m y^n + \overline{x^m y^n} | m \in \mathbb{Z}, n \geq 0\}, \\ \mathbb{C}_q^- &= \text{span}\{x^m y^n - \overline{x^m y^n} | m \in \mathbb{Z}, n > 0\}. \end{aligned}$$

Let M, N be two positive integers. We have a Lie superalgebra $gl(M, N)(\mathbb{C}_q)$ of $(M + N)$ by $(M + N)$ matrices with entries from \mathbb{C}_q .

We form a central extension of Lie superalgebra $gl(M, N)(\mathbb{C}_q)$ as was done in [G] and [CG].

$$(1.4) \quad \widehat{gl(M, N)}(\mathbb{C}_q) = gl(M, N)(\mathbb{C}_q) \oplus \left(\sum_{n \in \Lambda(q)} \oplus \mathbb{C}c(n) \right) \oplus \mathbb{C}c_y$$

with bracket

$$(1.5) \quad [A(x^m y^n), B(x^p y^s)]_s = A(x^m y^n)B(x^p y^s) - (-1)^{\deg A \deg B} B(x^p y^s)A(x^m y^n) \\ + mq^{np} \text{str}(AB) \delta_{m+p,0} \delta_{n+s,0} c(n+s) + nq^{np} \text{str}(AB) \delta_{m+p,0} \delta_{n+s,0} c_y$$

for $m, p, n, s \in \mathbb{Z}$, $A, B \in gl(M, N)_\alpha$, $\alpha = \bar{0}$ or $\bar{1}$, where str is the super-trace of the Lie superalgebra $gl(M, N)$, $c(u)$ with $u \in \Lambda(q)$ and c_y are central elements of $\widehat{gl(M, N)}(\mathbb{C}_q)$, \bar{t} means $\bar{t} \in \mathbb{Z}/\Lambda(q)$, for $t \in \mathbb{Z}$.

Now we present the examples of Lie superalgebra graded by the root system of type $B(0, N)$. We first set

$$J = \begin{pmatrix} 1 & 0 \\ 0 & -I_{2N} \end{pmatrix}, G = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & I_N \\ 0 & -I_N & 0 \end{pmatrix} \in M_{2N+1}(\mathbb{C}_q).$$

Then, G and J are invertible $(2N+1) \times (2N+1)$ -matrices. Using the matrix G and J , we define a superspace \mathcal{S} with:

$$\mathcal{S}_{\bar{0}} = \{X \in gl(1, 2N)(\mathbb{C}_q)_{\bar{0}} | \bar{X}^t G + GX = 0\}$$

$$\mathcal{S}_{\bar{1}} = \{X \in gl(1, 2N)(\mathbb{C}_q)_{\bar{1}} | \bar{X}^t G - JGX = 0\}$$

We can easily see that \mathcal{S} is a subalgebra of $gl(1, 2N)(\mathbb{C}_q)$ over \mathbb{C} . The general form of a matrix in \mathcal{S} is

$$(1.6) \quad \begin{pmatrix} a & b & c \\ \bar{c}^t & A & S \\ -\bar{b}^t & T & -\bar{A}^t \end{pmatrix} \quad \text{with } \bar{a} = -a \quad \bar{S}^t = S \quad \text{and} \quad \bar{T}^t = T,$$

where A, S, T are $N \times N$ sub-matrices. Then the Lie superalgebra $\mathcal{G} = [\mathcal{S}, \mathcal{S}]_s$, is a $B(0, N)$ -graded Lie superalgebra.

As in [AABGP], we easily know that:

$$\mathcal{G} = \{Y \in gl(1, 2N)(\mathbb{C}_q) | \text{str}(Y) \equiv 0 \bmod [\mathbb{C}_q, \mathbb{C}_q]\}.$$

Putting

$$(1.7) \quad \mathcal{H} = \left\{ \sum_{i=1}^N a_i (e_{ii} - e_{N+i, N+i}) | a_i \in \mathbb{C} \right\},$$

then \mathcal{H} is a N -dimensional abelian subalgebra of \mathcal{G} . Defining $\delta_i \in \mathcal{H}^*, i = 1, \dots, N$, by

$$(1.8) \quad \delta_i \left(\sum_{j=1}^N a_j (e_{jj} - e_{N+j, N+j}) \right) = a_i$$

for $i = 1, \dots, N$. Setting $\mathcal{G}_\alpha = \{x \in \mathcal{G} | [h, x]_s = \alpha(h)x, \text{ for all } h \in \mathcal{H}\}$ as usual, we have

$$(1.9) \quad \mathcal{G} = \mathcal{G}_0 \oplus \sum_{i \neq j} \mathcal{G}_{\delta_i - \delta_j} \oplus \sum_{i < j} (\mathcal{G}_{\delta_i + \delta_j} \oplus \mathcal{G}_{-\delta_i - \delta_j}) \oplus \sum_i (\mathcal{G}_{\delta_i} \oplus \mathcal{G}_{-\delta_i} \oplus \mathcal{G}_{2\delta_i} \oplus \mathcal{G}_{-2\delta_i}),$$

where

$$(1.10) \quad \begin{aligned} \mathcal{G}_{\delta_i - \delta_j} &= \text{span}_{\mathbb{C}} \{ \tilde{f}_{ij}(m, n) = x^m y^n e_{ij} - \overline{x^m y^n} e_{N+j, N+i} | m, n \in \mathbb{Z} \}, \\ \mathcal{G}_{\delta_i + \delta_j} &= \text{span}_{\mathbb{C}} \{ \tilde{g}_{ij}(m, n) = x^m y^n e_{i, N+j} + \overline{x^m y^n} e_{j, N+i} | m, n \in \mathbb{Z} \}, \\ \mathcal{G}_{-\delta_i - \delta_j} &= \text{span}_{\mathbb{C}} \{ \tilde{h}_{ij}(m, n) = -x^m y^n e_{N+i, j} - \overline{x^m y^n} e_{N+j, i} | m, n \in \mathbb{Z} \}, \\ \mathcal{G}_{2\delta_i} &= \text{span}_{\mathbb{C}} \{ \tilde{g}_{ii}(m, n) = (x^m y^n + \overline{x^m y^n}) e_{i, N+i} | m, n \in \mathbb{Z} \}, \\ \mathcal{G}_{-2\delta_i} &= \text{span}_{\mathbb{C}} \{ \tilde{h}_{ii}(m, n) = -(x^m y^n + \overline{x^m y^n}) e_{N+i, i} | m, n \in \mathbb{Z} \}, \\ \mathcal{G}_{\delta_i} &= \text{span}_{\mathbb{C}} \{ \tilde{e}_i(m, n) = -x^m y^n e_{i, 0} - \overline{x^m y^n} e_{0, N+i} | m, n \in \mathbb{Z} \}, \\ \mathcal{G}_{-\delta_i} &= \text{span}_{\mathbb{C}} \{ \tilde{e}_i^*(m, n) = x^m y^n e_{N+i, 0} - \overline{x^m y^n} e_{0, i} | m, n \in \mathbb{Z} \} \end{aligned}$$

and

$$\mathcal{G}_0 = \text{span}_{\mathbb{C}} \{ \tilde{f}_{ii}(m, n) + \tilde{e}_0(m, n), \tilde{e}_0(p, s) | 1 \leq i \leq N, m, n \in \mathbb{Z}, p \notin \Lambda(q) \text{ or } s \notin \Lambda(q) \}$$

where $\tilde{e}_0(m, n) = -(x^m y^n - \overline{x^m y^n}) e_{0, 0}$.

We then have a central extension of \mathcal{G}

$$(1.11) \quad \widehat{\mathcal{G}} = \mathcal{G} \oplus \left(\sum_{n \in \Lambda(q)} \oplus \mathbb{C} c(n) \right) \oplus \mathbb{C} c_y$$

with bracket as (1.5).

We have

Proposition 1.1

$$(1.12) \quad [\tilde{g}_{ij}(m, n), \tilde{g}_{kl}(p, s)]_s = 0$$

$$(1.13) \quad [\tilde{g}_{ij}(m, n), \tilde{f}_{kl}(p, s)]_s = -\delta_{il}q^{ms}\tilde{g}_{kj}(m+p, n+s) - \delta_{jl}q^{(s-n)m}\tilde{g}_{ki}(m+p, s-n)$$

$$(1.14) \quad \begin{aligned} & [\tilde{g}_{ij}(m, n), \tilde{h}_{kl}(p, s)]_s \\ = & -\delta_{ik}q^{-n(m+p)}\tilde{f}_{jl}(m+p, s-n) - \delta_{jk}q^{np}\tilde{f}_{il}(m+p, n+s) \\ & -\delta_{il}q^{-(mn+np+ps)}\tilde{f}_{jk}(m+p, -(n+s)) - \delta_{jl}q^{(n-s)p}\tilde{f}_{ik}(m+p, n-s) \\ & +mq^{np}\delta_{jk}\delta_{il}\delta_{m+p,0}\delta_{n+s,\bar{0}}(c(n+s) + c(-n-s)) \\ & +m\delta_{ik}\delta_{jl}\delta_{m+p,0}\delta_{n-s,\bar{0}}(c(n-s) + c(s-n)) \end{aligned}$$

$$(1.15) \quad [\tilde{g}_{ij}(m, n), \tilde{e}_k(p, s)]_s = [\tilde{g}_{ij}(m, n), \tilde{e}_0(p, s)]_s = 0$$

$$(1.16) \quad [\tilde{g}_{ij}(m, n), \tilde{e}_k^*(p, s)]_s = -\delta_{ik}q^{-n(m+p)}\tilde{e}_j(m+p, s-n) - \delta_{jk}q^{np}\tilde{e}_i(m+p, n+s)$$

$$(1.17) \quad [\tilde{g}_{ij}(m, n), \tilde{e}_0(p, s)]_s = 0$$

$$(1.18) \quad \begin{aligned} [\tilde{f}_{ij}(m, n), \tilde{f}_{kl}(p, s)]_s &= \delta_{jk}q^{np}\tilde{f}_{il}(m+p, n+s) - \delta_{il}q^{sm}\tilde{f}_{kj}(m+p, n+s) \\ &- 2mq^{np}\delta_{jk}\delta_{il}\delta_{m+p,0}\delta_{n+s,\bar{0}}c(n+s) \end{aligned}$$

$$(1.19) \quad [\tilde{f}_{ij}(m, n), \tilde{h}_{kl}(p, s)]_s = -\delta_{ik}q^{-n(m+p)}\tilde{h}_{jl}(m+p, s-n) - \delta_{il}q^{ms}\tilde{h}_{kj}(m+p, n+s)$$

$$(1.20) \quad [\tilde{f}_{ij}(m, n), \tilde{e}_k(p, s)]_s = \delta_{jk}q^{np}\tilde{e}_i(m+p, n+s)$$

$$(1.21) \quad [\tilde{f}_{ij}(m, n), \tilde{e}_k^*(p, s)]_s = -\delta_{ik}q^{-n(m+p)}\tilde{e}_j^*(m+p, s-n)$$

$$(1.22) \quad [\tilde{f}_{ij}(m, n), \tilde{e}_0(p, s)]_s = 0$$

$$(1.23) \quad [\tilde{h}_{ij}(m, n), \tilde{h}_{kl}(p, s)]_s = 0$$

$$(1.24) \quad [\tilde{h}_{ij}(m, n), \tilde{e}_k(p, s)]_s = \delta_{jk} q^{np} \tilde{e}_i^*(m + p, n + s) + \delta_{ik} q^{-n(m+p)} \tilde{e}_j^*(m + p, s - n)$$

$$(1.25) \quad [\tilde{h}_{ij}(m, n), \tilde{e}_k^*(p, s)]_s = 0$$

$$(1.26) \quad [\tilde{h}_{ij}(m, n), \tilde{e}_0(p, s)]_s = 0$$

$$(1.27) \quad [\tilde{e}_i(m, n), \tilde{e}_k(p, s)]_s = q^{m(s-n)} \tilde{g}_{ki}(m + p, s - n)$$

$$(1.28) \quad \begin{aligned} & [\tilde{e}_i(m, n), \tilde{e}_k^*(p, s)]_s \\ &= \delta_{ik} q^{-n(m+p)} \tilde{e}_0(m + p, s - n) + q^{p(n-s)} \tilde{f}_{ik}(m + p, n - s) \\ & \quad - m \delta_{ik} \delta_{m+p,0} \delta_{\overline{n-s}, \overline{0}} (c(n - s) + c(s - n)) \end{aligned}$$

$$(1.29) \quad [\tilde{e}_i(m, n), \tilde{e}_0(p, s)]_s = -q^{np} \tilde{e}_i(m + p, n + s) + q^{p(n-s)} \tilde{e}_i(m + p, n - s)$$

$$(1.30) \quad [\tilde{e}_i^*(m, n), \tilde{e}_k^*(p, s)]_s = q^{m(s-n)} \tilde{h}_{ki}(m + p, s - n)$$

$$(1.31) \quad [\tilde{e}_i^*(m, n), \tilde{e}_0(p, s)]_s = -q^{np} \tilde{e}_i^*(m + p, n + s) + q^{p(n-s)} \tilde{e}_i^*(m + p, n - s)$$

$$(1.32) \quad \begin{aligned} & [\tilde{e}_0(m, n), \tilde{e}_0(p, s)]_s \\ &= -(q^{np} - q^{sm}) \tilde{e}_0(m + p, n + s) - (q^{m(s-n)} - q^{-n(m+p)}) \tilde{e}_0(m + p, s - n) \\ & \quad + m q^{np} \delta_{m+p,0} \delta_{\overline{n+s}, \overline{0}} (c(n + s) + c(-n - s)) \\ & \quad - m \delta_{m+p,0} \delta_{\overline{n-s}, \overline{0}} (c(n - s) + c(s - n)) \end{aligned}$$

for all $m, p, n, s \in \mathbb{Z}$ and $1 \leq i, j, k, l \leq N$.

Remark 1.1 For our situation in the decomposition of $B(0, N)$ -graded Lie superalgebras in §5[Be2], $A = \mathbb{C}_q^+$ and $B = \mathbb{C}_q^-$.

2 The module construction

In this section, we follow the method in [G] and [CG] to construct representations for the Lie superalgebras which are given in Section 1. The idea goes back to [FF].

Let \mathcal{R} be an associative algebra. Let $\rho = \pm 1$. We define a ρ -bracket on \mathcal{R} as follow:

$$(2.1) \quad \{a, b\}_\rho = ab + \rho ba, \quad a, b \in \mathcal{R}.$$

It is easy to see that

$$(2.2) \quad \{a, b\}_\rho = \rho \{b, a\}_\rho$$

for $a, b, c \in \mathcal{R}$.

Define \mathfrak{a} to be the unital associative algebra with $2N$ generators $a_i, a_i^*, 1 \leq i \leq N$, subject to relations

$$(2.3) \quad \{a_i, a_j\}_- = \{a_i^*, a_j^*\}_- = 0, \quad \text{and} \quad \{a_i, a_j^*\}_- = -\delta_{ij}.$$

Let the associative algebra $\alpha(N, -)$ be generated by

$$(2.4) \quad \{u(m) | u \in \bigoplus_{i=1}^N (\mathbb{C}a_i \oplus \mathbb{C}a_i^*), m \in \mathbb{Z}\}$$

with the relations

$$(2.5) \quad \{u(m), v(n)\}_- = \{u, v\}_- \delta_{m+n, 0}.$$

The normal ordering is defined as in [FF](see also [F2] or [CG]).

$$(2.6) \quad \begin{aligned} : u(m)v(n) : &= \begin{cases} u(m)v(n) & \text{if } n > m, \\ \frac{1}{2}(u(m)v(n) + v(n)u(m)) & \text{if } n = m, \\ v(n)u(m) & \text{if } n < m, \end{cases} \\ &= : v(n)u(m) : \end{aligned}$$

for $n, m \in \mathbb{Z}, u, v \in \mathfrak{a}$.

Next we consider an extension of the algebra $\alpha(N, -)$. The generators

$$(2.7) \quad \{e(m) | m \in \mathbb{Z}\}$$

span an infinite-dimensional Clifford algebra with relations

$$(2.8) \quad \{e(m), e(n)\}_+ = e(m)e(n) + e(n)e(m) = -\delta_{n+m,0}.$$

Let $\alpha_\tau(N)$ denote the algebra obtained by adjoining to $\alpha(N, -)$ the generators (2.7) with relations (2.8) and

$$(2.9) \quad \{a_i(m), e(n)\}_\tau = 0 = \{a_i^*(m), e(n)\}_\tau, \text{ for } \tau = \pm 1.$$

The normal ordering is given as follows

$$(2.10) \quad \begin{aligned} :e(m)e(n): &:= \begin{cases} e(m)e(n) & \text{if } n > m, \\ \frac{1}{2}(e(m)e(n) - e(n)e(m)) & \text{if } n = m, \\ -e(n)e(m) & \text{if } n < m, \end{cases} \\ :a_i(m)e(n): &:= a_i(m)e(n) = -\tau e(n)a_i(m), \\ :a_i^*(m)e(n): &:= a_i^*(m)e(n) = -\tau e(n)a_i^*(m), \end{aligned}$$

for $n, m \in \mathbb{Z}, 1 \leq i, j \leq N$. Set

$$(2.11) \quad \theta(n) = \begin{cases} 1, & \text{for } n > 0, \\ \frac{1}{2}, & \text{for } n = 0, \\ 0, & \text{for } n < 0, \end{cases} \quad \text{then } 1 - \theta(n) = \theta(-n).$$

We have

$$(2.12) \quad \begin{aligned} :a_i(m)a_j(n): &:= a_i(m)a_j(n) = a_j(n)a_i(m), \\ :a_i^*(m)a_j^*(n): &:= a_i^*(m)a_j^*(n) = a_j^*(n)a_i^*(m), \end{aligned}$$

and

$$(2.13) \quad \begin{aligned} a_i(m)a_j^*(n) &:= a_i(m)a_j^*(n) : -\delta_{ij}\delta_{m+n,0}\theta(m-n), \\ a_j^*(n)a_i(m) &:= a_i(m)a_j^*(n) : +\delta_{ij}\delta_{m+n,0}\theta(n-m), \\ e(m)e(n) &:= e(m)e(n) : -\delta_{m+n,0}\theta(m-n). \end{aligned}$$

Let $\alpha(N, -)^+$ be the subalgebra generated by $a_i(n), a_j^*(m), a_k^*(0)$, for $n, m > 0$, and $1 \leq i, j, k \leq N$. Let $\alpha(N, -)^-$ be the subalgebra generated by $a_i(n), a_j^*(m), a_k(0)$, for $n, m < 0$, and $1 \leq i, j, k \leq N$. Those generators in $\alpha(N, -)^+$ are called annihilation operators while those in $\alpha(N, -)^-$ are called creation operators. Let $V(N, -)$ be a simple $\alpha(N, -)$ -module containing an

element v_0 , called a “vacuum vector”, and satisfying

$$(2.14) \quad \alpha(N, -)^+ v_0 = 0.$$

So all annihilation operators kill v_0 and

$$(2.15) \quad V(N, -) = \alpha(N, -)^- v_0.$$

Let V_0 be a simple Clifford module for the Clifford algebra generated by (2.7) with relations (2.8) and containing “vacuum vector” v'_0 , which is killed by annihilation operators. (Here we call $e(m)$ annihilation operator if $m > 0$, or a creation operator if $m < 0$. $e(0)$ acts as scalar.) Because of (2.9), we see that the $\alpha_\tau(N)$ -module

$$(2.16) \quad V_\tau(N) = V(N, -) \otimes V_0 = \alpha_\tau(N) v'_0$$

is simple.

Now we define our operators on $V_\tau(N)$. For any $m, n \in \mathbb{Z}$, $1 \leq i, j \leq N$, set

$$(2.17) \quad f_{ij}(m, n) = \sum_{s \in \mathbb{Z}} q^{-ns} : a_i(m-s) a_j^*(s) :$$

$$(2.18) \quad g_{ij}(m, n) = \sum_{s \in \mathbb{Z}} q^{-ns} : a_i(m-s) a_j(s) :$$

$$(2.19) \quad h_{ij}(m, n) = \sum_{s \in \mathbb{Z}} q^{-ns} : a_i^*(m-s) a_j^*(s) :$$

$$(2.20) \quad e_i(m, n) = \sum_{s \in \mathbb{Z}} q^{-ns} : a_i(m-s) e(s) :$$

$$(2.21) \quad e_i^*(m, n) = \sum_{s \in \mathbb{Z}} q^{-ns} : a_i^*(m-s) e(s) :$$

$$(2.22) \quad e_0(m, n) = \sum_{s \in \mathbb{Z}} q^{-ns} : e(m-s) e(s) :.$$

We will list only those commutation relations involving $e(s)$ (See [CG]).

Lemma 2.1 *We have*

$$(2.23) \quad [a_i(m) a_j(n), a_k(p) e(s)] = [a_i(m) a_j(n), e(p) e(s)] = 0$$

$$(2.24) \quad [a_i(m) a_j(n), a_k^*(p) e(s)] = -\delta_{ik} \delta_{m+p,0} a_j(n) e(s) - \delta_{jk} \delta_{n+p,0} a_i(m) e(s)$$

$$(2.25) \quad [a_i(m) a_j^*(n), a_k(p) e(s)] = \delta_{jk} \delta_{n+p,0} a_i(m) e(s),$$

$$(2.26) \quad [a_i(m)a_j^*(n), a_k^*(p)e(s)] = -\delta_{ik}\delta_{m+p,0}a_j^*(n)e(s),$$

$$(2.27) \quad [a_i(m)a_j^*(n), e(p)e(s)] = [a_i^*(m)a_j^*(n), a_k^*(p)a_l^*(s)] = 0$$

$$(2.28) \quad [a_i^*(m)a_j^*(n), a_k(p)e(s)] = \delta_{jk}\delta_{n+p,0}a_i^*(m)e(s) + \delta_{ik}\delta_{m+p,0}a_j^*(m)e(s),$$

$$(2.29) \quad [a_i^*(m)a_j^*(n), a_k^*(p)e(s)] = [a_i^*(m)a_j^*(n), e(p)e(s)] = 0$$

$$(2.30) \quad \{a_i(m)e(n), a_k(p)e(s)\}_+ = \tau\delta_{n+s,0}a_i(m)a_k(p)$$

$$(2.31) \quad \{a_i(m)e(n), a_k^*(p)e(s)\}_+ = \tau\delta_{n+s,0}a_k^*(p)a_i(m) + \tau\delta_{ik}\delta_{m+p,0}e(n)e(s)$$

$$(2.32) \quad [a_i(m)e(n), e(p)e(s)] = \delta_{n+s,0}a_i(m)e(p) - \delta_{n+p,0}a_i(m)e(s)$$

$$(2.33) \quad \{a_i^*(m)e(n), a_k^*(p)e(s)\}_+ = \tau\delta_{n+s,0}a_i^*(m)a_k^*(p)$$

$$(2.34) \quad [a_i^*(m)e(n), e(p)e(s)] = \delta_{n+s,0}a_i^*(m)e(p) - \delta_{n+p,0}a_i^*(m)e(s)$$

$$(2.35) \quad [e(m)e(n), e(p)e(s)] \\ = -\delta_{n+p,0}e(m)e(s) + \delta_{m+p,0}e(n)e(s) - \delta_{n+s,0}e(p)e(m) + \delta_{m+s,0}e(p)e(n)$$

for $m, n, p, s \in \mathbb{Z}$ and $1 \leq i, j, k, l \leq N$.

Proof We only check some of them.

$$\begin{aligned}
& \{a_i(m)e(n), a_k(p)e(s)\}_+ = a_i(m)e(n)a_k(p)e(s) + a_k(p)e(s)a_i(m)e(n) \\
&= -\tau(a_i(m)a_k(p)e(n)e(s) + a_k(p)a_i(m)e(s)e(n)) \\
&= -\tau(a_i(m)a_k(p)(e(n)e(s) + e(s)e(n))) \\
&= \tau\delta_{n+s,0}a_i(m)a_k(p);
\end{aligned}$$

$$\begin{aligned}
& \{a_i(m)e(n), a_k^*(p)e(s)\}_+ = a_i(m)e(n)a_k^*(p)e(s) + a_k^*(p)e(s)a_i(m)e(n) \\
&= -\tau(a_i(m)a_k^*(p)e(n)e(s) + a_k^*(p)a_i(m)e(s)e(n)) \\
&= -\tau(a_k^*(p)a_i(m) - \delta_{ik}\delta_{m+p,0})e(n)e(s) - \tau a_k^*(p)a_i(m)e(s)e(n) \\
&= \tau\delta_{n+s,0}a_k^*(p)a_i(m) + \tau\delta_{ik}\delta_{m+p,0}e(n)e(s);
\end{aligned}$$

$$\begin{aligned}
& [a_i(m)e(n), e(p)e(s)] = a_i(m)e(n)e(p)e(s) - e(p)e(s)a_i(m)e(n) \\
&= a_i(m)(e(n)e(p)e(s) - (-\tau)^2e(p)e(s)e(n)) \\
&= a_i(m)(-\delta_{n+p,0}e(s) - e(p)e(n)e(s) - e(p)e(s)e(n)) \\
&= a_i(m)(\delta_{n+s,0}e(p) - \delta_{n+p,0}e(s)) \\
&= \delta_{n+s,0}a_i(m)e(p) - \delta_{n+p,0}a_i(m)e(s);
\end{aligned}$$

$$\begin{aligned}
& [e(m)e(n), e(p)e(s)] = e(m)e(n)e(p)e(s) - e(p)e(s)e(m)e(n) \\
&= e(m)(-\delta_{n+p,0} - e(p)e(n))e(s) - e(p)e(s)e(m)e(n) \\
&= -\delta_{n+p,0}e(m)e(s) - e(m)e(p)e(n)e(s) - e(p)e(s)e(m)e(n) \\
&= -\delta_{n+p,0}e(m)e(s) - (-\delta_{m+p,0} - e(p)e(m))e(n)e(s) - e(p)e(s)e(m)e(n) \\
&= -\delta_{n+p,0}e(m)e(s) + \delta_{m+p,0}e(n)e(s) + e(p)(e(m)e(n)e(s) - e(s)e(m)e(n)) \\
&= -\delta_{n+p,0}e(m)e(s) + \delta_{m+p,0}e(n)e(s) + e(p)(\delta_{m+s,0}e(n) - \delta_{n+s,0}e(m)) \\
&= -\delta_{n+p,0}e(m)e(s) + \delta_{m+p,0}e(n)e(s) - \delta_{n+s,0}e(p)e(m) + \delta_{m+s,0}e(p)e(n).
\end{aligned}$$

So (2.30), (2.31), (2.32) and (2.35) hold true. ■

In what follows we shall mean $\frac{q^{mn}-1}{q^n-1} = m$ if $n \in \Lambda(q)$. This will make our formula more concise.

Next we list all brackets that are needed. For all $m, p, n, s \in \mathbb{Z}$ and $1 \leq i, j, k, l \leq N$,

Proposition 2.1

$$[g_{ij}(m, n), g_{kl}(p, s)] = 0$$

Proposition 2.2

$$[g_{ij}(m, n), f_{kl}(p, s)] = -\delta_{il}q^{ms}g_{kj}(m + p, n + s) - \delta_{jl}q^{(s-n)m}g_{ki}(m + p, s - n)$$

Proposition 2.3

$$\begin{aligned} & [g_{ij}(m, n), h_{kl}(p, s)] \\ = & -\delta_{ik}q^{-n(m+p)}f_{jl}(m + p, s - n) - \delta_{jk}q^{np}f_{il}(m + p, n + s) \\ & -\delta_{il}q^{-(mn+np+ps)}f_{jk}(m + p, -(n + s)) - \delta_{jl}q^{(n-s)p}f_{ik}(m + p, n - s) \\ & +\delta_{ik}\delta_{jl}\delta_{m+p,0}\frac{1}{2}(q^{s-n} + 1)\frac{q^{m(s-n)} - 1}{q^{s-n} - 1} + \delta_{jk}\delta_{il}\delta_{m+p,0}q^{np}\frac{1}{2}(q^{s+n} + 1)\frac{q^{m(s+n)} - 1}{q^{s+n} - 1} \end{aligned}$$

Proposition 2.4

$$[g_{ij}(m, n), e_k(p, s)] = [g_{ij}(m, n), e_0(p, s)] = 0$$

$$[g_{ij}(m, n), e_k^*(p, s)] = -\delta_{ik}q^{-n(m+p)}e_j(m + p, s - n) - \delta_{jk}q^{np}e_i(m + p, n + s)$$

Proposition 2.5

$$\begin{aligned} [f_{ij}(m, n), f_{kl}(p, s)] &= \delta_{jk}q^{np}f_{il}(m + p, n + s) - \delta_{il}q^{sm}f_{kj}(m + p, n + s) \\ &\quad -\delta_{jk}\delta_{il}q^{np}\delta_{m+p,0}\frac{1}{2}(q^{s+n} + 1)\frac{q^{m(s+n)} - 1}{q^{s+n} - 1} \end{aligned}$$

Proposition 2.6

$$[f_{ij}(m, n), h_{kl}(p, s)] = -\delta_{ik}q^{-n(m+p)}h_{jl}(m + p, s - n) - \delta_{il}q^{ms}h_{kj}(m + p, n + s)$$

Proposition 2.7

$$\begin{aligned} [f_{ij}(m, n), e_k(p, s)] &= \delta_{jk}q^{np}e_i(m + p, n + s) \\ [f_{ij}(m, n), e_k^*(p, s)] &= -\delta_{ik}q^{-n(m+p)}e_j^*(m + p, s - n) \\ [f_{ij}(m, n), e_0(p, s)] &= 0 \end{aligned}$$

Proposition 2.8

$$[h_{ij}(m, n), h_{kl}(p, s)] = 0$$

Proposition 2.9

$$[h_{ij}(m, n), e_k(p, s)] = \delta_{jk} q^{np} e_i^*(m + p, n + s) + \delta_{ik} q^{-n(m+p)} e_j^*(m + p, s - n)$$

$$[h_{ij}(m, n), e_k^*(p, s)] = [h_{ij}(m, n), e_0(p, s)] = 0$$

Proposition 2.10

$$\{e_i(m, n), e_k(p, s)\}_+ = \tau q^{m(s-n)} g_{ki}(m + p, s - n)$$

$$\{e_i(m, n), e_k^*(p, s)\}_+ = \tau \delta_{ik} q^{-n(m+p)} e_0(m + p, s - n) + \tau q^{p(n-s)} f_{ik}(m + p, n - s)$$

$$- \tau \delta_{ik} \delta_{m+p,0} \frac{1}{2} (q^{s-n} + 1) \frac{q^{m(s-n)} - 1}{q^{s-n} - 1}$$

$$[e_i(m, n), e_0(p, s)] = -q^{np} e_i(m + p, n + s) + q^{p(n-s)} e_i(m + p, n - s)$$

Proposition 2.11

$$\{e_i^*(m, n), e_k^*(p, s)\}_+ = \tau q^{m(s-n)} h_{ki}(m + p, s - n)$$

$$[e_i^*(m, n), e_0(p, s)] = -q^{np} e_i^*(m + p, n + s) + q^{p(n-s)} e_i^*(m + p, n - s)$$

Proposition 2.12

$$[e_0(m, n), e_0(p, s)]$$

$$= -(q^{np} - q^{sm}) e_0(m + p, n + s) + \delta_{m+p,0} q^{np} \frac{1}{2} (q^{n+s} + 1) \frac{q^{m(n+s)} - 1}{q^{n+s} - 1}$$

$$- (q^{m(s-n)} - q^{-n(m+p)}) e_0(m + p, s - n) - \delta_{m+p,0} \frac{1}{2} (q^{s-n} + 1) \frac{q^{m(s-n)} - 1}{q^{s-n} - 1}$$

We only give proofs for 2.10 and 2.12. The proof for the others is either similar or easy.

Proofs of 2.10 and 2.12:

Note that from [CG], we have

$$(2.36) \quad \sum_{t \in \mathbb{Z}} q^{-xt} \left(\theta(-2t) - \theta(-2m - 2t) \right) = \frac{1}{2} (q^x + 1) \frac{q^{mx} - 1}{q^x - 1}.$$

By (2.30),(2.31),(2.32) and (2.35), we have

$$(2.37) \quad \{e_i(m, n), a_k(p)e(s)\}_+ = \tau q^{ns} a_k(p) a_i(m + s),$$

$$(2.38) \quad \{e_i(m, n), a_k^*(p)e(s)\}_+ = \tau q^{ns} a_k^*(p) a_i(m + s) + \tau \delta_{ik} q^{-n(m+p)} e(m + p) e(s),$$

$$(2.39) \quad [e_i(m, n), e(p)e(s)] = q^{ns} a_i(m + s) e(p) - q^{np} a_i(m + p) e(s),$$

and

$$(2.40) \quad [e_0(m, n), e(p)e(s)] = -(q^{np} - q^{-n(m+p)}) e(m + p) e(s) - (q^{ns} - q^{-n(m+s)}) e(p) e(m + s),$$

so we get

$$\begin{aligned} & \{e_i(m, n), q^{-st} : a_k(p - t) e(t) : \}_+ = \tau q^{nt-st} a_k(p - t) a_i(m + t) \\ &= \tau q^{m(s-n)} q^{-(s-n)(m+t)} a_k(p - t) a_i(m + t) \\ &= \tau q^{m(s-n)} q^{-(s-n)(m+t)} : a_k(p - t) a_i(m + t) :, \end{aligned}$$

$$\begin{aligned} & \{e_i(m, n), q^{-st} : a_k^*(p - t) e(t) : \}_+ \\ &= q^{-st} \left(\tau \delta_{ik} q^{-n(m+p-t)} e(m + p - t) e(t) + \tau q^{nt} a_k^*(p - t) a_i(m + t) \right) \\ &= \tau \delta_{ik} q^{-n(m+p)} q^{-(s-n)t} e(m + p - t) e(t) + \tau q^{-(s-n)t} a_k^*(p - t) a_i(m + t) \\ &= \tau \delta_{ik} q^{-n(m+p)} q^{-(s-n)t} \left(: e(m + p - t) e(t) : - \delta_{m+p,0} \theta(m + p - 2t) \right) \\ & \quad \tau q^{-(s-n)t} \left(: a_i(m + t) a_k^*(p - t) : + \delta_{ik} \delta_{m+p,0} \theta(p - m - 2t) \right) \\ &= \tau \delta_{ik} q^{-n(m+p)} q^{-(s-n)t} : e(m + p - t) e(t) : + \tau q^{p(n-s)} q^{-(n-s)(p-t)} : a_i(m + t) a_k^*(p - t) : \\ & \quad - \tau \delta_{ik} \delta_{m+p,0} q^{-(s-n)t} (\theta(-2t) - \theta(-2m - 2t)); \end{aligned}$$

$$\begin{aligned} & [e_i(m, n), q^{-st} : e(p - t) e(t) :] \\ &= q^{-st} (q^{n(p-t)} a_i(m + p - t) e(t) + q^{nt} a_i(m + t) e(p - t)) \\ &= -q^{np} q^{-(n+s)t} : a_i(m + p - t) e(t) : + q^{p(n-s)} q^{-(n-s)(p-t)} : a_i(m + t) e(p - t) : \end{aligned}$$

and

$$\begin{aligned}
& [e_0(m, n), q^{-st} : e(p-t)e(t) :] \\
&= -q^{-st} \left((q^{n(p-t)} - q^{-n(m+p-t)})e(m+p-t)e(t) - (q^{nt} - q^{-n(m+t)})e(p-t)e(m+t) \right) \\
&= -q^{-st} (q^{n(p-t)} - q^{-n(m+p-t)}) (: e(m+p-t)e(t) : -\delta_{m+p,0}\theta(m+p-2t)) \\
&\quad -q^{-st} (q^{nt} - q^{-n(m+t)}) (: e(p-t)e(m+t) : -\delta_{m+p,0}\theta(p-m-2t)) \\
&= -q^{np}q^{-(s+n)t} : e(m+p-t)e(t) : +q^{-n(m+p)}q^{-(s-n)t} : e(m+p-t)e(t) : \\
&\quad -q^{m(s-n)}q^{-(s-t)(m+t)} : e(p-t)e(m+t) : +q^{sm}q^{-(n+s)(m+t)} : e(p-t)e(m+t) : \\
&\quad +\delta_{m+p,0}q^{np}q^{-(n+s)t}(\theta(-2t) - \theta(-2m-2t)) \\
&\quad -\delta_{m+p,0}q^{-(s-n)t}(\theta(-2t) - \theta(-2m-2t))
\end{aligned}$$

by (2.36), we see that Proposition 2.10 and 2.12 hold true. \blacksquare

To find the correspondence of the homomorphism, we need to modify our operators by shifting some central elements.

For Proposition 2.3, we see that, if $n+s \in \Lambda(q)$ and $n-s \in \Lambda(q)$,

$$\begin{aligned}
& [g_{ij}(m, n), h_{kl}(p, s)] \\
&= -\delta_{ik}q^{-n(m+p)}f_{jl}(m+p, s-n) - \delta_{jk}q^{np}f_{il}(m+p, n+s) \\
&\quad -\delta_{il}q^{-(mn+np+ps)}f_{jk}(m+p, -(n+s)) - \delta_{jl}q^{(n-s)p}f_{ik}(m+p, n-s) \\
&\quad +\delta_{ik}\delta_{jl}\delta_{m+p,0}m + \delta_{jk}\delta_{il}\delta_{m+p,0}q^{np}m.
\end{aligned}$$

If $n+s \in \mathbb{Z} \setminus \Lambda(q)$ and $n-s \in \Lambda(q)$, then

$$\begin{aligned}
& [g_{ij}(m, n), h_{kl}(p, s)] \\
&= -\delta_{ik}q^{-n(m+p)}f_{jl}(m+p, s-n) - \delta_{jl}q^{(n-s)p}f_{ik}(m+p, n-s) \\
&\quad -\delta_{jk}q^{np} \left(f_{il}(m+p, n+s) + \frac{1}{2}\delta_{il}\delta_{m+p,0}\frac{q^{n+s}+1}{q^{n+s}-1} \right) \\
&\quad -\delta_{il}q^{-(mn+np+ps)} \left(f_{jk}(m+p, -n-s) + \frac{1}{2}\delta_{jk}\delta_{m+p,0}\frac{q^{-n-s}+1}{q^{-n-s}-1} \right) \\
&\quad +\delta_{ik}\delta_{jl}\delta_{m+p,0}m.
\end{aligned}$$

Similarly, if $n + s \in \Lambda(q)$ and $n - s \in \mathbb{Z} \setminus \Lambda(q)$

$$\begin{aligned}
& [g_{ij}(m, n), h_{kl}(p, s)] \\
= & -\delta_{jk}q^{np}f_{il}(m + p, n + s) - \delta_{il}q^{-(mn+np+ps)}f_{jk}(m + p, -n - s) \\
& -\delta_{ik}q^{-n(m+p)}\left(f_{jl}(m + p, s - n) + \frac{1}{2}\delta_{jl}\delta_{m+p,0}\frac{q^{s-n}+1}{q^{s-n}-1}\right) \\
& -\delta_{jl}q^{(n-s)p}\left(f_{ik}(m + p, n - s) + \frac{1}{2}\delta_{ik}\delta_{m+p,0}\frac{q^{n-s}+1}{q^{n-s}-1}\right) \\
& +\delta_{jk}\delta_{il}\delta_{m+p,0}q^{np}m.
\end{aligned}$$

By the above two relations, we have if $n + s, n - s \in \mathbb{Z} \setminus \Lambda(q)$

$$\begin{aligned}
& [g_{ij}(m, n), h_{kl}(p, s)] \\
= & -\delta_{jk}q^{np}\left(f_{il}(m + p, n + s) + \frac{1}{2}\delta_{il}\delta_{m+p,0}\frac{q^{n+s}+1}{q^{n+s}-1}\right) \\
& -\delta_{il}q^{-(mn+np+ps)}\left(f_{jk}(m + p, -n - s) + \frac{1}{2}\delta_{jk}\delta_{m+p,0}\frac{q^{-n-s}+1}{q^{-n-s}-1}\right) \\
& -\delta_{ik}q^{-n(m+p)}\left(f_{jl}(m + p, s - n) + \frac{1}{2}\delta_{jl}\delta_{m+p,0}\frac{q^{s-n}+1}{q^{s-n}-1}\right) \\
& -\delta_{jl}q^{(n-s)p}\left(f_{ik}(m + p, n - s) + \frac{1}{2}\delta_{ik}\delta_{m+p,0}\frac{q^{n-s}+1}{q^{n-s}-1}\right)
\end{aligned}$$

Using the same method, for Proposition 2.5 we have, if $n + s \in \Lambda(q)$,

$$\begin{aligned}
& [f_{ij}(m, n), f_{kl}(p, s)] \\
= & \delta_{jk}q^{np}f_{il}(m + p, n + s) - \delta_{il}q^{sm}f_{kj}(m + p, n + s) - \delta_{jk}\delta_{il}q^{np}\delta_{m+p,0}m.
\end{aligned}$$

If $n + s \in \mathbb{Z} \setminus \Lambda(q)$, then

$$\begin{aligned}
& [f_{ij}(m, n), f_{kl}(p, s)] \\
= & \delta_{jk}q^{np}\left(f_{il}(m + p, n + s) + \frac{1}{2}\delta_{il}\delta_{m+p,0}\frac{q^{n+s}+1}{q^{n+s}-1}\right) \\
& -\delta_{il}q^{sm}\left(f_{kj}(m + p, n + s) + \frac{1}{2}\delta_{jk}\delta_{m+p,0}\frac{q^{n+s}+1}{q^{n+s}-1}\right).
\end{aligned}$$

For Proposition 2.10, if $n - s \in \Lambda(q)$,

$$\begin{aligned}
& \{e_i(m, n), e_k^*(p, s)\}_+ \\
= & \tau\delta_{ik}q^{-n(m+p)}e_0(m + p, s - n) + \tau q^{p(n-s)}f_{ik}(m + p, n - s) - \tau\delta_{ik}\delta_{m+p,0}m.
\end{aligned}$$

If $n + s \in \mathbb{Z} \setminus \Lambda(q)$, then

$$\begin{aligned} & \{e_i(m, n), e_k^*(p, s)\}_+ \\ = & \tau \delta_{ik} q^{-n(m+p)} \left(e_0(m+p, s-n) + \frac{1}{2} \delta_{m+p,0} \frac{q^{s-n} + 1}{q^{s-n} - 1} \right) \\ & + \tau q^{p(n-s)} \left(f_{ik}(m+p, n-s) + \frac{1}{2} \delta_{jk} \delta_{m+p,0} \frac{q^{n-s} + 1}{q^{n-s} - 1} \right). \end{aligned}$$

For Proposition 2.12, if $n + s \in \Lambda(q)$ and $n - s \in \Lambda(q)$,

$$\begin{aligned} & [e_0(m, n), e_0(p, s)] \\ = & -(q^{np} - q^{sm})e_0(m+p, n+s) - (q^{m(s-n)} - q^{-n(m+p)})e_0(m+p, s-n) \\ & + \delta_{m+p,0} q^{np} m - \delta_{m+p,0} m. \end{aligned}$$

If $n + s \in \mathbb{Z} \setminus \Lambda(q)$ and $n - s \in \Lambda(q)$, then

$$\begin{aligned} & [e_0(m, n), e_0(p, s)] \\ = & -(q^{np} - q^{sm}) \left(e_0(m+p, n+s) + \frac{1}{2} \delta_{m+p,0} \frac{q^{n+s} + 1}{q^{n+s} - 1} \right) \\ & - (q^{m(s-n)} - q^{-n(m+p)})e_0(m+p, s-n) - \delta_{m+p,0} m. \end{aligned}$$

If $n + s \in \Lambda(q)$ and $n - s \in \mathbb{Z} \setminus \Lambda(q)$

$$\begin{aligned} & [e_0(m, n), e_0(p, s)] \\ = & -(q^{np} - q^{sm})e_0(m+p, n+s) + \delta_{m+p,0} q^{np} m \\ & - (q^{m(s-n)} - q^{-n(m+p)}) \left(e_0(m+p, s-n) + \frac{1}{2} \delta_{m+p,0} \frac{q^{s-n} + 1}{q^{s-n} - 1} \right). \end{aligned}$$

If $n + s, n - s \in \mathbb{Z} \setminus \Lambda(q)$

$$\begin{aligned} & [e_0(m, n), e_0(p, s)] \\ = & -(q^{np} - q^{sm}) \left(e_0(m+p, n+s) + \frac{1}{2} \delta_{m+p,0} \frac{q^{n+s} + 1}{q^{n+s} - 1} \right) \\ & - (q^{m(s-n)} - q^{-n(m+p)}) \left(e_0(m+p, s-n) + \frac{1}{2} \delta_{m+p,0} \frac{q^{s-n} + 1}{q^{s-n} - 1} \right). \end{aligned}$$

Therefore, if we define

$$\begin{aligned}
(2.41) \quad F_{ij}(m, n) &= \begin{cases} f_{ij}(m, n), & \text{for } n \in \Lambda(q) \\ f_{ij}(m, n) + \frac{1}{2}\delta_{ij}\delta_{m,0}\frac{q^n+1}{q^n-1}, & \text{for } n \in \mathbb{Z} \setminus \Lambda(q) \end{cases} \\
G_{ij}(m, n) &= g_{ij}(m, n), \quad H_{ij}(m, n) = h_{ij}(m, n), \\
E_i(m, n) &= e_i(m, n), \quad E_i^*(m, n) = e_i^*(m, n), \\
E_0(m, n) &= \begin{cases} e_0(m, n), & \text{for } n \in \Lambda(q) \\ e_0(m, n) + \frac{1}{2}\delta_{m,0}\frac{q^n+1}{q^n-1}, & \text{for } n \in \mathbb{Z} \setminus \Lambda(q) \end{cases}
\end{aligned}$$

we have

Theorem 2.1 $V_\tau(N)$ is a module for the Lie superalgebra $\widehat{\mathcal{G}}$ under the action given by (for $\tau = \pm 1$)

$$\begin{aligned}
\pi(\tilde{g}_{ij}(m, n)) &= \tau G_{ij}(m, n), & \pi(\tilde{f}_{ij}(m, n)) &= F_{ij}(m, n), \\
\pi(\tilde{h}_{ij}(m, n)) &= \tau H_{ij}(m, n), & \pi(\tilde{e}_i(m, n)) &= \tau E_i(m, n), \\
\pi(\tilde{e}_i^*(m, n)) &= E_i^*(m, n), & \pi(\tilde{e}_0(m, n)) &= E_0(m, n), \\
\pi(c_x) &= -\frac{1}{2}, & \pi(c_y) &= 0.
\end{aligned}$$

References

- [AABGP] B.N. Allison, S. Azam, S. Berman, Y. Gao and A. Pianzola, Extended affine Lie algebras and their root systems. Mem. Amer. Math. Soc. (605) **126** (1997)
- [ABG] B.N. Allison, G. Benkart, Y. Gao, Lie algebras graded by the root systems $BC_r, r \geq 2$. Mem. Amer. Math. Soc. (751) **158** (2002)
- [BE1] G. Benkart, A. Elduque, Lie superalgebras graded by the root system $A(m; n)$, J. Lie Theory **13** (2003) 387–400.
- [BE2] G. Benkart, A. Elduque, Lie superalgebras graded by the root system $B(m, n)$. Selecta Math. (N.S.) **9** (2003) 313–360.
- [BGK] S. Berman, Y. Gao and Y. Krylyuk, Quantum tori and the structure of elliptic quasi-simple Lie algebras, J. Funct. Anal. **135** (1996) 339C389.
- [CG] H. Chen, Y. Gao, BC_N -graded Lie algebras arising from fermionic representations, math.QA/0601284

- [F1] I.B. Frenkel, Spinor representations of affine Lie algebras, Proc. Nat. Acad. Sci. USA **77** (1980) 6303–6306.
- [F2] I.B. Frenkel, Two constructions of affine Lie algebra representations and Boson-Fermion correspondence in quantum field theory, J. Funct. Anal. **44** (1981) 259–327.
- [FF] A.J. Feingold and I.B. Frenkel, Classical affine algebras, Adv. Math. **56** (1985) 117–172.
- [G] Y. Gao, Fermionic and bosonic representations of the extended affine Lie algebra $\widetilde{gl_N(\mathbb{C}_q)}$, Canad. Math. Bull. **45** (2002) 623–633.
- [GN] E. García and E. Neher, Tits-Kantor-Koecher superalgebras of Jordan superpairs covered by grids. Comm. Algebra **31** (2003) 3335–3375.
- [K] V.G. Kac, Lie superalgebras, Adv. Math. **26** (1977) 8–96.
- [KP] V.G. Kac and D.H. Peterson, Spin and wedge representations of infinite-dimensional Lie algebras and groups, Proc. Nat. Acad. Sci. USA **78** (1981) 3308–3312.
- [M] Y.I. Manin, Topics in noncommutative geometry. Princeton University Press (1991).

Hongjia Chen, Shikui Shang
 Department of Mathematics
 University of Science and Technology of China
 Hefei, Anhui
 P. R. China 230026
 Email: hjchen@mail.ustc.edu.cn
 Email: skshang@mail.ustc.edu.cn

Yun Gao
 Department of Mathematics and Statistics
 York University
 Toronto, Ontario
 Canada M3J 1P3
 Email: ygao@yorku.ca